

Date:

Assignment No: 2: General science and ability

sir majid Raza chahbari

Name: Naveed Ali

ID # 32359 - OB.

Topic: Malaria and typhoid.

### MALARIA:-

Malaria is an infection of RBC's and liver.

Malaria is caused by plasmodium carried by female mosquito called anopheles. When the mosquito with plasmodium bites a healthy person it injects/release plasmodium in the blood of that person which then cause malarial disease in the body.

### CAUSES OF MALARIA :-

Malaria is caused if a mosquito infected with the plasmodium parasite bites somebody. There are four kinds of malaria parasites that can infect humans.

- (1) Plasmodium vivax.
- (2) Plasmodium ovale
- (3) Plasmodium malariae
- (4) Plasmodium falciparum

Plasmodium falciparum causes a more severe form of the disease and those who contract this form of malaria have a higher risk of death. An infected mother can also pass the disease to her baby at birth.

This is known as congenital malaria. Malaria can be transmitted by blood so it can be transmitted through an organ transplant, a transfusion and use of shared needles or syringes.

Date:

## SIGNS AND SYMPTOMS OF MALARIA

The patient of this disease experienced recurring attacks of chills and fevers. Sweating, headache, vomiting and diarrhoea. Malaria signs and symptoms typically began within few weeks after being bitten by infected mosquito. However, some types of malaria parasites can lie dormant in human body for 9 years.

## TREATMENT

Blood tests can show the presence of the parasites. and chloroquine and primaquine collectively used for the treatment of P. vivax, P. ovale and P. malaria.

The treatment for P. falciparum is chloroquine for sensitive areas and Quinine, sulfadoxine and pyrimethamine for quinine resistant areas.

## PREVENTION

Preventional measures for malaria are: sleeping under mosquito nets, stagnant water should be avoided, using insect repellents and cover the limbs.

## TYPHOID

It is bacterial infection that can lead to a high fever, diarrhea and vomiting. It is caused by the bacteria salmonella typhi which is a gram negative rod shaped flagellated bacterium whose only reservoir is the human body.

## TRANSMISSION

This disease can spread through contaminated food and drinking contaminated water. Washing fruits and vegetables with contaminated water can also spread typhoid fever.

Date: \_\_\_\_\_

it can also spread through flies.

### SIGNS AND SYMPTOMS :-

Signs and symptoms include

- 1) Fever that starts low and increased daily reaching as high as 104°F.
- 2) Headache
- 3) weakness and fatigue.
- 4) poor appetite
- 5) abdominal pain
- 6) Diarrhea or constipation.
- 7) Reddish spots on abdomen due to bleeding under abdominal skin.
- 8) Nausea or vomiting.
- 9) if left untreated intestinal ulcers may occurs.

### TREATMENT :-

it is a bacterial infection so antibiotics like ciprofloxacin are used to treat the infection. Antipyretic drugs are also given to normalize temperature & and complete bed rest is advised.

### PREVENTION :-

Preventional measures are:

- 1) Frequently washing hands
- 2) Avoiding drinking untreated water.
- 3) Avoid raw fruits and vegetables.
- 4) Choose hot foods.

