| | DATE | | |
|------------|---|--|--|
| | Prevention: | | |
| -7 | Sleep under insecticide hednet- | | |
| -7 | Full body clothing- | | |
| -7 | Mosquito repellents- | | |
| -> | Use of insecticide sprays | | |
| -7 | Full body clothing- Mosquito repellents- Use of insecticide sprays Early detection & treatment using antibiotics- | | |
| | | | |
| - | reatments: | | |
| | Following are the known treatments for modaria | | |
| | Microscopy:- | | |
| =7 | It is a gold standard method in diagnosting malaria- | | |
| =7 | In microscopy, three samples of thick & Ithin | | |
| | blood are looked under athe microscope- | | |
| | "Thick Sample — 7 For identifying the parasites Thin Sample — 7 For identifying the specis-" | | |
| | Thin Sample -> For identifying the specis-" | | |
| | | | |
| | Antimalarial Drugs:- | | |
| i-) | | | |
| ü-) | Chloroquine | | |
| <u>w</u>) | Quinine etc. | | |
| | | | |
| | | | |

Define madaia, Typhoid, It's symptoms, causes, preventions and treatments? DATE_ YPHOID:-Definition:-Typhoid Fever also known as Enteric Fever lize threatning injection caused the bacterium Salmonella Typhi-" symptoms:-Salmonella Typhi lives only in humans-=> Person affected by this disease carry bacteria their bloodstream intestinal tractand Prolonged high zever Muscle aches & Fatigue and Abdominal pain -Extreme headach => | Constipation and diarrhea-Causes:--7 The Ga Contaminated 700d and water is The Cause of malaria.

Another cause is the close contact with who is carrying salmonella bacteria - Urbanization and climate charge have t have the Potential to increase typhoid-Increasing resistance to autibiotic treatment making 7 easter for typhoid to spread-Use full Prevention: Good personal hygiene mainly is the of typhoid-Acess to safe water-Adequate Sanitation. Typoid Jaccination -

Question:

| | DATE | * |
|-------------|--|------------------------------|
| | Treatments:- | |
| | Antibiotics: | |
| 1-) | Fluoroquinolores - | |
| ù-) | Azithromycin- | |
| \ü-) | Ceftriaxone- | |
| (-vi | Carbapenems- | |
| | • 4 | |
| | Two additional vaccines have | been used to prevent typor - |
| Handard Co. | .Anactivate d | Live attenuated |
| =7 | An injectable vaccine-(IV injection) | => Given orany (capsule) |
| =) | Criven to people 2 years & above | -1 Given to 6 years and |
| =) | For travellers, it is reccommended | older- |
| | to give single dose 2 weeks | |
| | before travel- | Good answers!! |
| . =) | Repeated dose for people who remain at risk. | |
| | who remain at risk. | |
| | | |
| | | |
| | | |