Assignment #1 Hajra Sarfraz Batch 59 Rollno 32405 Topic: Polio: Symptoms, Causes, treatment & Prevention. Poliomyelitis: Poliomyelitis, most commonly known as polio, is an infectious disease caused by polio virus which affects the motor neurons in brain and spinal cord. It can cause paralysis of legs and arms, and if it affects the muscles of diaphragm it may cause death. Symptoms: 90-95% people infected by people does not show symptoms. 5-10%. people show mild symptoms like fatigue, fever, headache, throat pain abdominal pain nausea and vomiting. While 0.5% people show sever symptoms such as muscle weakness, paresthesia (feelings of pins & needles in the legs), paralysis and flobby limbs.

Causes:

Poliomyelitis is caused by Polio vinus which is a non-envelopeel \$5 RNA vinus. It has three scrotypes PV1,

PV2, PV3.

PV2 and PV3

are completely

eradicate, only PV1

exists now.

Capsid 3, RNA, RNA, RNA, RNA, Structure of Polio Vixus.

Mechanism of Action:- Polio virus enters into
the mouth and more down into throat
where it replicates. It mores down further
into stomach and intestine, where it makes
more and more cupies and then excrete
out of body in stool. If it succeeds
in entering into the blood, our immune
system fights with it and kills if
between
immunity is strong. This fight both virus
and immune cells cause the mild
symptoms. But if virus enter becomes
able to enter into central nervous system,

Keep the description of a single
heading brief and divide it into
itsubfleadings motor neurons in Brainstem
and spinal cord. It and damages the
a cell bodies of motor neurons in grey
matter. As the motor neurons damaged,
no electrical impulse goes to muscles and
activity of muscles reduced. The process is
rapid and cause complete paralysis of
muscles. If it affects the muscles of diaphingm,
breathing well be difficult, and patient
will died ultimately.




Diagram showing linkege between nervous system and muscles through motor neurons.

Made of Transmission: It is transmitted through two routes:

1 Fecal to oral route 1 Oral to oral route Diagnosis & Treatment: Diagnosis: Polio is diagnosed by symptoms and lab tests. There are three types of lab tests are done including throat culture, stool culture and cerbs spinal fluid culture. Treatment: There is not a specific treatment for polio. However, supportive care is provided to reduce the pain and Symptoms. Supportive care includes .. @ Flyids 1 Rest 3 Iron lung/ventillators (4) Dain medication invented Iron lung, was a device, in 20th century used for patients with damaged mudes of diaphragm. Iron lyng was a closed device which covers the toold whole body except face. It changes the pressure level and allow the diaphram to expand and keep breathing. Now, it has been replaced by ventilators.

9

visus and suffer from

Prevention:	
The ma	jor preventive measure
is vaccination which projects human from	
virus by inducing immonity. There are two	
bypes of vaccinations	available for polio:
O Sabin	o Salk.
Sabin	Salk
It uses live afterwated	It uses killed form
virus.	of virus
Its 3 doses give 95%	Its 1 dose provides 90%.
protection.	while 3 doses provide
	99 /. projection
It is oral vaccine	is intramuscular vaccine
and given orall	which is injected into
	muscles of leg or arm
It provides hymoral	It provides only
as well as cell -	hymoral immunity.
mediated immunity.	
1 out 2.5 million	The vivus connot be
people can get	visulent as it is
virulent form of	dead.

Other preventive measures: Pollo spread in greas of crowding, poor hygein and sanitation. That is why, we can prevent it - increasing sanitation standard - survellience. Polio Free Would The world aims to eradicate the polio and to turn into Polio-free world. By 1994, Averica were free of polio. Now America, Europe and westren Pacific completely eraplicated the virus. As reported on WHO in 2014 only 3 countries have polic cases including Pakistan, Afghanistan and Negeria By implementing the preventive measures, we can get rid of polio. Overall a good answer! But the answer is lengthy and will affect your time management. So

shorten it a bit