Mention the full qs statements for proper evaluation. Without that these are just notes and cannot be propagycevaluated Dengue is the most rapidly spreading mosquito-borne vival di se which couses sudden fever and a se pain in joints Symptoms: Nausea and vamiting Rashes on the skim Body arhe ethorgy and rest le 4-Sudden high fever Causal agent: Dengue wires is transmitted to humans through the bite of Aeder aegypti mosquido. General couses: When an infected Acdes mosquita bites a person it transmits dengte wires in that person if an uninfected magnito takes the blood of an injected person, it becomes infected. Now this masquito can transmit the virus to a healthy person.

Date	economia.	2274	and in	

Treatment:	: picolo M
Most cases of dengue fener	
at home with pain medicies	
minophen (paraceto mol). In	
italization is often needed.	
Prevention:	
Remouring stagment water	
Identification of dengue larvae	and larniciding .
Pluging water leakages	NAME OF THE PERSON OF THE PERS
Using mosquito nels and	squito repellents
blooring long-sleeped shirt	ts and parils.

6- Graving anti-dengue plants like lemon grass,

7- Spraying kerosene oil on the nearby water

bodies such as pools, puddles and ponds etc

marigald and mint etc.

Malaria: type of fever spread to human by some types of marquitar It is a life-threatening disease but Symptoms! High fener 2- Nousea and vointing 3. painful body neutranen Shinering of body due to Fatigue 6- Diarshoa Causal agent: This disease is coursed by a parasite called plasmodum, transmitted to humans by the bite of a female Anophelese nominit General Causes: when an infected me quite bites a human plasmodum en the bloodstream cousing malaria. When an uninfected masquita bites an injected person, it becomes injected and in

this way the disease is spread from person

-	-	-	
	ean		

1			
to person.	-4.0.9		
Treatment:			

Malaria is treated by using anti-malarial drugs.

Most common anti-marial drug is Chloroquine
phosphate.

## Prevention:

- 1. Using a parathyroid-containing insect spray in living and sleeping area during evening and night times
- 2. Using mosquita bed net
- 3- Using mosquito repellents.
- 4. Wearing clothes that cover most of the body
- 5- Dumping stagnant water
- 6. Speaying keron and on nearby water

Polio:
Polio, also known as Poliomyelitis, is a high
infectious disease caused by a virus. It as
attacks the nervous system and can cause
paralysis. Paralysis on affect arms, legs as
muscles that control breathing.
Symptoms:
1- Fatigue
2 Fener
3- Headache
4- Varieting and naura
5. Pain in limbs and stumach
6- Stiffness in the neck and sore throat
Casal agent:
palio is caused by palio wins It is very
contagious wires and can spread easily
from one person to another.
General Cosser:
Polio spreads through contact with infected
muces and phlegm through coughing an

beliefing ent: their tratmos descent stopping the frecer. For enample not washing hands after touching facces (like the zing diaper ar gaing belowing the ballroom) and in inspire (moralled and at by faceal material etc. Symptonics

Treatment:

There is no cure for polio.

Palie can be prevented by help of vaccine. which is given orally in form of drops. Every! child must receive the vaccine at both of? upto 6 neeks, and thereafter at 3 months, 45 months and 6 months of age. A booster is given at the age of 18 months more locus?

The selection of the se · market and a self about roomed and glassified General Course:

Typhoid: Typhoid fever is a bacterial infection that can spread throughout the body affecting many organs without proper treatment, it can be folal. Symptoms: Prolonged high fever Fatigue Headache 3-4-Nausea 5-Modominal pain Diarrhea or Constitució 6-Some people develop rash Causal agent: Typhoid is caused by bacterium called Salmonella Typhi. Once these barterio are injested, they multiply and stread int the blood stream: General Cause: It is usually spread through contaminated food or water.

Treatment:
Ardibictics are used to treat typhoid former.
Prevention:
1 Wash hands before eating or handling food
2- Bail maler before dienking: A illingett -0
3- Eat thoroughly cooked food
4- Dispose of gorsage properly
5- Get vaccinated.
6- Ensure proper sandation and sewage disposal
in the surroundings.
- and water buspared the character of the college of the
Hepatilis:
This disease is the inflamation of liver coursed
by a wird infection.
Symptons:
1- Yellowing of skin and whites of the eyes.
2- Dark virine or year wrine
3- Abdominal pain : 8 altrogott -d
4- Muscle ache
5. Fatique

	Date
	: meatinger
6. Lass caf appetite	and the state of the state of
7. Lethorgy	
mild fener	: mailingung
9- Pale foeces	and decay
Types of Hepatitis:	island Line
a- Hepatitis A:	midble those is
This type derives from an infection	visite of the second
hopatitis A views (HAV).	i (Brouming )
. It is most commonly transmitted by fa	oces from
food and walls contaminated of the	
a person infected with reporting A	1 universal
is a small unemeloped symmetri	COL RNA MOUN
which shares many of the charac	toxiation of
- the picomavirus family	Masmas of
. Its incubation period is about	
There is no specific treatment	for nepaires
5- Hepatitis B:	Server State of the Control of the C
. This type desines I som an infec	
hepatitis B ruins (HBV).	Supplied &

ill is most commonly transmitted through contact
with infected body fluid as sharing razors and
syringes with an injected person its
Hepatilis & wins, a rembor of hepanda wins
group, double stronded ONA views which replicate.
Ms incubation period is variable with range 1 do
6 months.
Antiviral medication is used for treatment.
C. Hepatitis C
· This type comes from hepatitis C visus (HCV).
ell is transmitted through direct contact with.
infected body fluids.
· Hepotitis C is an er relaped, single-stranded
RNA Marine Local Company of the Comment of
· Antificial medication is used to treat chronic hepatitis
C.
d. 1-lepatitis D:
ell is also called della hebatitis
HOV is transmitted thrown contact with wounds
or contact with injected blood
· Hepatitis D wows is single-stranded circular RNA.
. 15 1

Date				
	•	٠	٠	٠

chronic hepatitis o may be treated by medication called interferons. e- Hepatitis Englished dur monde e it is a water-borne disease caused by hepatitis E wins. (HEV) habranda alduch . It is found mostly in areas with poor sanitation. . It is non-em sped, single-stranded RNA www. · There is no specific medication to treat 2 rititoglaH -Prevention: · Use of vaccines of HAV and HBV · Screening blood used for transfusion · Safe injection produces · Using hygier cally fit food and water

· Maintain proper sanitation and cleanliness of . surroundings.